


=> d 11
 L1 HAS NO ANSWERS
 L1 STR



REP G1=(0-3) CH
 VAR G2=2/1
 NODE ATTRIBUTES:
 CHARGE IS E+1 AT 3
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC 1
 NUMBER OF NODES IS 13

STEREO ATTRIBUTES: NONE

=> fil caplus
 COST IN U.S. DOLLARS
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
0.49	0.71

FILE 'CAPLUS' ENTERED AT 09:30:26 ON 02 MAR 2010
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 2 Mar 2010 VOL 152 ISS 10
 FILE LAST UPDATED: 1 Mar 2010 (20100301/ED)
 REVISED CLASS FIELDS (/NCL) LAST RELOADED: Dec 2009
 USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Dec 2009

CAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 12

L3 46 L2

=> s 13 and py<=2003

24050416 PY<=2003

L4 12 L3 AND PY<=2003

=> d bib abs 1-12

L4 ANSWER 1 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2003:837081 CAPLUS

DN 139:337885

TI Preparation of acyloxypyrrolidinium salts as M3 muscarinic antagonists

IN Prat Quinones, Maria; Fernandez Forner, Maria Dolors

PA Almirall Prodesfarma S.A., Spain

SO PCT Int. Appl., 72 pp.

CODEN: PIXXD2

DT Patent

LA English




FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003087094	A2	20031023	WO 2003-EP3786	20030411 <--
	WO 2003087094	A3	20040318		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	ES 2206021	A1	20040501	ES 2002-889	20020416
	ES 2206021	B1	20050801		
	CA 2482536	A1	20031023	CA 2003-2482536	20030411 <--
	AU 2003233967	A1	20031027	AU 2003-233967	20030411 <--
	AU 2003233967	B2	20090806		
	EP 1497284	A2	20050119	EP 2003-727294	20030411
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	BR 2003009167	A	20050125	BR 2003-9167	20030411
	NZ 535884	A	20071130	NZ 2003-535884	20030411
	RU 2320657	C2	20080327	RU 2004-133371	20030411
	MX 2004010076	A	20041213	MX 2004-10076	20041013
	IN 2004DN03157	A	20070112	IN 2004-DN3157	20041013
	ZA 2004008335	A	20051102	ZA 2004-8335	20041014
	NO 2004004826	A	20050114	NO 2004-4826	20041105
	US 20050282875	A1	20051222	US 2005-510680	20050720
	US 7192978	B2	20070320		
	US 20070129420	A1	20070607	US 2007-648581	20070103
PRAI	ES 2002-889	A	20020416		
	WO 2003-EP3786	W	20030411		
	US 2005-510680	A3	20050720		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 139:337885

GI



AB Pyrrolidinium derivs. I [R = (un)substituted Ph, naphthalenyl, 5,6,7,8-tetrahydronaphthalenyl, benzo[1,3]dioxolyl, biphenyl, heteroarom.; R1 = alkyl; R2 = CR3R4R5, Q; R3 = 2-furyl, 3-furyl, 2-thienyl, 3-thienyl; R4 = 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, cycloalkyl; R5 = H, OH, Me, CH2OH; Q1 = CH2, CH2CH2, O, OCH2, S, SCH2, CH:CH; A = (un)substituted CH:CH, CH2, CO, O, S, S(O), SO2, NH; m = 0-8; n = 0-4] were prepared for use in therapy as antagonists of M3 muscarinic receptors (no data). Thus, (3R)-3-pyrrolidinol was treated with 2-(3-bromopropyl)thiophene to give (3R)-1-(3-thien-2-ylpropyl)pyrrolidinol which was treated with Me 2-hydroxy-2,2-dithen-2-ylacetate and quaternized to give the pyrrolidinium salt II.

OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2002:889558 CAPLUS

DN 137:369966

TI Preparation of enantiomerically pure basic [(cyclopentyl- or cyclohexylhydroxyphenylacetyl)oxy]-1,1-dimethylpyrrolidinium salts, their muscarinic receptor binding affinity, and use as treatment for obstructive respiratory disease

IN Noe, Christian; Mutschler, Ernst; Lambrecht, Gunter; Elgert, Michael; Elgert, Ruth Irene; Czeche, Sittah; Waelbroeck, Magali

PA Germany

SO U.S. Pat. Appl. Publ., 19 pp., Cont.-in-part of U.S. 6,307,060. CODEN: USXXCO

DT Patent

LA English


FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20020173536	A1	20021121	US 2001-901217	20010709 <--
	US 6613795	B2	20030902		
	WO 9821183	A1	19980522	WO 1997-AT245	19971111 <--
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
	RW:	GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
	EP 1369414	A1	20031210	EP 2003-5233	19971111 <--
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, AL			
	EP 1371645	A1	20031217	EP 2003-5232	19971111 <--
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, AL			
	US 6307060	B1	20011023	US 1999-309960	19990511 <--
	US 20030220400	A1	20031127	US 2003-601542	20030623 <--
	US 7253182	B2	20070807		
PRAI	AT 1996-1973	A	19961111		
	WO 1997-AT245	W	19971111		
	US 1999-309960	A2	19990511		
	EP 1997-911049	A3	19971111		
	US 2001-901217	A3	20010709		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 137:369966

GI



AB Disclosed are enantiomerically pure cyclic aminoalc. esters of arylcycloalkylhydroxycarboxylic acids with at least 90% enantiomeric excess of the (3R,2'R), (3S,2'R), (3R,2'S), or (3S,2'S) configured enantiomer. Thus, [(cyclopentyl (or cyclohexyl) hydroxyphenylacetyl)oxyl] pyrrolidinium salts I (R = cyclopentyl, cyclohexyl, X = bromide, iodide, fluoride, chloride) were prepared by reacting (3R) or (3S)-1-methyl-3-pyrrolidinol with the corresponding phenylacetate, followed by preparation of the tartrate intermediates and quaternization. Inhalable powder and aerosol formulations of the compds. were also prepared. The muscarinic binding affinity of I were examined using rabbit vas deferens, guinea pic atrium, guinea pig ileum, and human M1, M2, and M3 receptors.


OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L4 ANSWER 3 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2002:154498 CAPLUS

DN 137:15684

TI Studies on a soft glycopyrrolate analog, SG-1
 AU Ji, F.; Wu, W.-M.; Bodor, N.
 CS Center for Drug Discovery, University of Florida, Gainesville, FL, USA
 SO Pharmazie (2002), 57(2), 138-141
 CODEN: PHARAT; ISSN: 0031-7144
 PB Govi-Verlag Pharmazeutischer Verlag GmbH
 DT Journal
 LA English
 GI



AB A short-acting soft drug analog (I) of glycopyrrolate (G) was developed by retrometabolic design in order to minimize systemic side effects and optimize the therapeutic index. I was synthesized by: (a) esterification of phenylacetic acid with N-methyl-3-pyrrolidinol by DCC to obtain N-methyl-3-pyrrolidinyl phenylacetate; (b) reaction of lithium salt of above phenylacetates with cyclopentanone to obtain N-methyl-3-pyrrolidinyl 3-(1'-hydroxycyclopentyl)phenylacetate; and (c) quaternization with Me bromoacetate in acetonitrile to give the designed product. To evaluate the pharmacol. effect of I, its mydriatic activity in rabbit eyes was compared to that of glycopyrrolate. At the pharmacodynamically equivalent doses (the lowest dose that induces the maximum response) of I (1%) and glycopyrrolate (0.1%), the mydriatic activities lasted for 5 and 100 h, resp. Compared to glycopyrrolate, the intrinsic pupil dilation potency of I was lower (.apprx.1/10th) but the duration of action was much shorter (<1/20th) as I is susceptible to facile enzymic hydrolysis/deactivation in the rabbit eyes. In vitro metabolism and stability investigations further supported this finding. In vitro half lives of I in rat plasma, blood, and 20% liver and lung tissue homogenates were 15.62, 53.86, 263.43, and 318.35 min, resp. In human plasma and blood, half-lives were 19.93 and 88.32 min, resp. I was relatively stable under acidic conditions (pH 5 and lower). I is a promising, clin. useful short acting anticholinergic.

OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2002:78639 CAPLUS

DN 137:194970

TI Receptor binding studies of soft anticholinergic agents

AU Huang, Fenglei; Buchwald, Peter; Browne, Clinton E.; Farag, Hassan H.; Wu, Whei-Mei; Ji, Fubao; Hochhaus, Guenther; Bodor, Nicholas

CS Center for Drug Discovery, College of Pharmacy, University of Florida, Gainesville, FL, 32610-0497, USA

SO PharmSci [online computer file] (2001), 3(4), No pp. given

CODEN: PHARFY; ISSN: 1522-1059

URL: http://www.pharmsci.org/scientificjournals/pharmsci/journal/pdf/01_30.pdf

PB American Association of Pharmaceutical Scientists
 DT Journal; (online computer file)
 LA English
 OS CASREACT 137:194970
 AB Receptor binding studies were performed on 24 soft anticholinergic agents and 5 conventional anticholinergic agents using 4 cloned human muscarinic receptor subtypes. The measured pKi values of the soft anticholinergic agents ranged from 6.5 to 9.5, with the majority being in the range of 7.5 to 8.5. Strong correlation was observed between the pKis determined here and the

p42 values measured earlier in guinea pig ileum contraction assays. The corresponding correlation coeffs. (r²) were 0.80, 0.73, 0.81, and 0.78 for pKi(m1), pKi(m2), pKi(m3), and pKi(m4), resp. Quant. structure-activity relationship (QSAR) studies were also performed, and good characterization could be obtained for the soft anticholinergics containing at least 1 tropine moiety in their structure. For these compds., the potency as measured by the pKi values was found to be related to geometric, electronic, and lipophilicity descriptors. A linear regression equation using ovality (Oe), dipole moment (D), and a calculated log octanol-water partition coefficient (QLogP) gave reasonably good descriptions (r = 0.88) for the pKi(m3) values.

OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)
 RE.CNT 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN
 AN 2002:51415 CAPLUS
 DN 136:118468
 TI Preparation of 2-aryl-2-hydroxyacetic acid ester derivatives as muscarinic M3 receptor antagonists
 IN Ogino, Yoshio; Kurihara, Hideki; Matsuda, Kenji; Numazawa, Tomoshige; Otake, Norikazu; Noguchi, Kazuhito
 PA Banyu Pharmaceutical Co., Ltd., Japan
 SO PCT Int. Appl., 138 pp.
 CODEN: PIXXD2
 DT Patent
 LA Japanese
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002004402	A1	20020117	WO 2001-JP5987	20010710 <--
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 2001071027	A	20020121	AU 2001-71027	20010710 <--
	CA 2415468	A1	20030110	CA 2001-2415468	20010710 <--
	EP 1302458	A1	20030416	EP 2001-949925	20010710 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	AU 2001271027	B2	20050707	AU 2001-271027	20010710
	US 20030191316	A1	20031009	US 2003-332617	20030110 <--
	US 6846835	B2	20050125		
	US 20050065211	A1	20050324	US 2004-983613	20041109
	US 7192969	B2	20070320		

	US 20070129397	A1	20070607	US 2007-648614	20070103
	US 7504432	B2	20090317		
PRAI	JP 2000-210591	A	20000711		
	WO 2001-JP5987	W	20010710		
	US 2003-332617	A3	20030110		
	US 2004-983613	A3	20041109		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 136:118468

AB Compds. of the general formula $\text{ArC(OH)(R1)CO}_2\text{A}$ [wherein A is a group of the general formula $-\text{B1-N+R2R3R4.X-}$ or $-\text{B2-NR5CR6:NR7}$; Ar is aryl or heteroaryl, any of which may be substituted; B1 and B2 are each an aliphatic hydrocarbon group; R1 is fluorinated cycloalkyl; R2, R3 and R4 are each lower alkyl, or a single bond or alkylene, any of which is bonded to B1, or alternatively R2 and R3 may be united to form alkylene; R5 and R7 are each hydrogen, lower alkyl, or a single bond or alkylene, any of which is bonded to B2; R6 is hydrogen, lower alkyl, or N(R8)R9 ; R8 and R9 are independently hydrogen or lower alkyl; and X- is an anion] are prepared. These compds. exhibit selective muscarinic M3 receptor antagonism with little side effects and are suitable for administration by inhalation and useful as therapeutic agents for respiratory system diseases including chronic obstructive pulmonary diseases, chronic bronchitis, asthma, chronic airway obstruction, pulmonary fibrosis, pulmonary emphysema, or rhinitis. Thus, reductive methylation of piperidin-4-yl (2R)-((1R)-3,3-difluoropentyl)-2-hydroxy-2-phenylethanoate by formaldehyde and sodium cyanoborohydride in the presence of ZnCl_2 in MeOH at room

temperature

for 30 min gave 1-methylpiperidin-4-yl (2R)-((1R)-3,3-difluoropentyl)-2-hydroxy-2-phenylethanoate which was quaternized by Me bromide in MeCN at room temperature for 15 h to give 4-[[[(2R)-2-((1R)-3,3-difluorocyclopentyl)-2-hydroxy-2-phenylethanoyl]oxy]-1,1-dimethylpiperidinium bromide (I). In a muscarinic receptor M2 and M3 antagonism assay, 4-(((2R)-2-((1R)-3,3-difluorocyclopentyl)-2-hydroxy-2-phenylethanoyl)oxy)-1,1-dimethylpiperidinium bromide in vitro exhibited KB of 9.6 nM for inhibiting the carbachol-induced reduction in heart beat in rat right atrium (muscarinic receptor M2 receptor) and that of 0.004 nM for inhibiting the carbachol-induced contraction of trachea (muscarinic receptor M3 receptor) with M2/M3 receptor ratio of 218. An ampule or a powder inhalation formulation containing I were described.

OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 2000:192082 CAPLUS

DN 133:758

TI Design, synthesis, and pharmacological evaluation of soft glycopyrrolate and its analog

AU Ji, F.; Huang, F.; Juhasz, A.; Wu, W.; Bodor, N.

CS Center for Drug Discovery, College of Pharmacy, University of Florida, Gainesville, FL, USA

SO Pharmazie (2000), 55(3), 187-191

CODEN: PHARAT; ISSN: 0031-7144

PB Govi-Verlag Pharmazeutischer Verlag

DT Journal

LA English

AB Glycopyrrolate is a quaternary anticholinergic drug. Like for other anticholinergics, the usefulness of this agent is limited by its side effects. In this study, based on the structure of glycopyrrolate, we designed a soft drug, methoxycarbonylphenylcyclopentylacetoxymethyl-N,N-dimethyl-3-pyrrolidinium Me sulfate (SG), and its analog, methoxycarbonylphenylcyclopentylacetoxymethyl-N,N,N-trimethylammonium Me

sulfate (SGA). These soft drugs are expected to be locally active, but systemically inactive in order to increase therapeutic index. SG and SGA were synthesized by (i) carboxylation of Me phenylcyclopentylacetate, (ii) esterification with N-methyl-3-pyrrolidinol (for SG) or 2-chloro-N,N-dimethylaminoethane (for SGA), and (iii) quaternization with di-Me sulfate. Receptor binding studies demonstrate that SG has muscarinic subtype selectivity (m3/m2). Guinea pig ileum pA2 assay indicates that activity of SG is moderate, and SG is about ten times more potent than SGA. The in vivo characterization of SG and SGA, both in mydriasis tests and in prevention of carbachol induced bradycardia, supported its soft nature. Applying SG or SGA into rabbit eyes, the dilation of the contralateral (water-treated) pupils was not observed. Glycopyrrolate application, however, caused dilation of the contralateral pupil, indicating a systemic effect of this drug. Cardiac studies were carried out by evaluating the protective effect of soft anticholinergics against carbachol induced bradycardia. The results indicate that SG and SGA were as potent as atropine-MeBr in preventing carbachol induced bradycardia in the rat; however, their durations of action were significantly shorter. In conclusion, the newly synthesized SG and SGA showed soft nature in the body. They are anticholinergics with subtype selectivity and moderate potency, and can be used as topical antiperspirants.

OSC.G 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)
 RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:341543 CAPLUS

DN 129:27887

OREF 129:5943a,5946a

TI Preparation of 1,1-dialkylpyrrolidinium-3-yl α -cycloalkylmandelate diastereomers and analogs as muscarinic M3 receptor ligands

IN Noe, Christian R.; Mutschler, Ernst; Lambrecht, Gunter; Elgert, Michael; Czeche, Sittah; Waelbroeck, Magali

PA Germany

SO PCT Int. Appl., 37 pp.

CODEN: PIXXD2

DT Patent

LA German

FAN.CNT 2


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9821183	A1	19980522	WO 1997-AT245	19971111 <--
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
	RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	CA 2271276	A1	19980522	CA 1997-2271276	19971111 <--
	CA 2271276	C	20090113		
	AU 9748560	A	19980603	AU 1997-48560	19971111 <--
	AU 745331	B2	20020321		
	EP 937041	A1	19990825	EP 1997-911049	19971111 <--
	EP 937041	B1	20030423		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	CN 1237159	A	19991201	CN 1997-199621	19971111 <--
	CN 100391942	C	20080604		

HU 9903791	A2	20000328	HU 1999-3791	19971111 <--
HU 9903791	A3	20011029		
NZ 336202	A	20001027	NZ 1997-336202	19971111 <--
JP 2001504459	T	20010403	JP 1998-521942	19971111 <--
AT 238280	T	20030515	AT 1997-911049	19971111 <--
PT 937041	E	20030930	PT 1997-911049	19971111 <--
ES 2195121	T3	20031201	ES 1997-911049	19971111 <--
EP 1369414	A1	20031210	EP 2003-5233	19971111 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, AL				
EP 1371645	A1	20031217	EP 2003-5232	19971111 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, AL				
RU 2238936	C2	20041027	RU 1999-112115	19971111
PL 195520	B1	20070928	PL 1997-332595	19971111
NO 9901056	A	19990511	NO 1999-1056	19990303 <--
NO 314354	B1	20030310		
US 6307060	B1	20011023	US 1999-309960	19990511 <--
US 20020173536	A1	20021121	US 2001-901217	20010709 <--
US 6613795	B2	20030902		
PRAI AT 1996-1973	A	19961111		
EP 1997-911049	A	19971111		
WO 1997-AT245	W	19971111		
US 1999-309960	A2	19990511		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 129:27887

GI



AB Title compds. [I; R = R1R4C(OH)CO2; R1 = (un)substituted cycloalkyl; R4 = (hetero)aryl; Z = NR2R3A; A = pharmacol. acceptable acid conjugate base; R2,R3 = (halo)alk(en)yl, (halo)alkynyl; Z1 = (CH2)1-3] were prepared Thus, (S)-1-methyl-3-pyrrolidinol was esterified by PhCR1(OH)CO2Me (R1 = cyclopentyl) and the resolved product quaternized to give (S)-I [R = (S)-PhCR1(OH)CO2, R1 = cyclopentyl, Z = NMe2I, Z1 = CH2]. Data for biol. activity of I were given.

OSC.G 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 8 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:58955 CAPLUS

DN 128:132422

OREF 128:25915a,25918a

TI Methods and compositions for treating urinary incontinence using enantiomerically enriched (SR)-glycopyrrolate

IN Fabiano, Vincent L.; McCullough, John R.

PA Sepracor, Inc., USA

SO PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9800109	A1	19980108	WO 1997-US11639	19970627 <--
	W: US				
	RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP	909163	A1	19990421	EP 1997-931548	19970627 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
PRAI	US 1996-21156P	P	19960701		
	WO 1997-US11639	W	19970627		
AB	A method for treating urinary incontinence, such as incontinence resulting from bladder detrusor muscle instability, using enantiomerically enriched (SR)-glycopyrrolate (I). The method comprises administering a therapeutically effective amount of enantiomerically enriched I, or a pharmaceutically acceptable salt thereof, substantially free of the (R,S)-enantiomer. Pharmaceutical compns. for the treatment of urinary incontinence comprising enantiomerically enriched I, or a pharmaceutically acceptable salt thereof, and an acceptable carrier are also disclosed. The antimuscarinic, spasmolytic, and Ca entry blocking effects of models of receptor binding and bladder function were studied for I.				
OSC.G	1	THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)			
RE.CNT	2	THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT			

L4 ANSWER 9 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:55524 CAPLUS

DN 128:119670

OREF 128:23363a,23366a

TI Methods and compositions for treating urinary incontinence using enantiomerically enriched (SS)-glycopyrrolate

IN Fabiano, Vincent L.; McCullough, John R.

PA Sepracor, Inc., USA

SO PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9800133	A1	19980108	WO 1997-US11645	19970627 <--
	W: US				
	RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP	920313	A1	19990609	EP 1997-932472	19970627 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	US 6063808	A	20000516	US 1998-214169	19981229 <--
PRAI	US 1996-21020P	P	19960701		
	WO 1997-US11645	W	19970627		
AB	A method for treating urinary incontinence, such as incontinence resulting from bladder detrusor muscle instability, using enantiomerically enriched (S,S)-glycopyrrolate (I). The method comprises administering a therapeutically effective amount of enantiomerically enriched I, or a pharmaceutically acceptable salt thereof, substantially free of the (R,R)-glycopyrrolate enantiomer. Pharmaceutical compns. for the treatment of urinary incontinence comprising enantiomerically enriched I or a pharmaceutically acceptable salt thereof, and an acceptable carrier are also disclosed. Pharmacol. data are given for binding to muscarinic receptor subtypes, Ca channels, and antimuscarinic-antispasmodic activity.				
OSC.G	2	THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)			
RE.CNT	1	THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT			

L4 ANSWER 10 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1998:55485 CAPLUS

DN 128:136512

OREF 128:26699a,26702a

TI Methods and compositions for treating urinary incontinence using enantiomerically enriched (R,R)-glycopyrrolate

IN Fabiano, Vincent L.; McCullough, John R.

PA Sepracor, Inc., USA

SO PCT Int. Appl., 29 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 9800016	A1	19980108	WO 1997-US11644	19970627 <--
	W: US				
	RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP	932401	A1	19990804	EP 1997-931551	19970627 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	US 6204285	B1	20010320	US 1998-214168	19981229 <--
PRAI	US 1996-20947P	P	19960701		
	WO 1997-US11644	W	19970627		

AB A method is disclosed for treating urinary incontinence, e.g. incontinence resulting from bladder detrusor muscle instability, using enantiomerically enriched (R, R)-glycopyrrolate. The method comprises administering a therapeutically effective amount of enantiomerically enriched (R, R)-glycopyrrolate, or a pharmaceutically acceptable salt thereof, substantially free of the (S, S)-glycopyrrolate enantiomer. Pharmaceutical compns. for the treatment of urinary incontinence, comprising enantiomerically enriched (R, R)-glycopyrrolate, or a pharmaceutically acceptable salt thereof, and an acceptable carrier, are also disclosed.

OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 11 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1965:54408 CAPLUS

DN 62:54408

OREF 62:9664f-g

TI The antiperspirant action of topically applied anticholinergics

AU MacMillan, F. S. Kilmer; Reller, Herbert H.; Synder, Fred H.

CS Procter & Gamble Co., Cincinnati, OH

SO Journal of Investigative Dermatology (1964), 43, 363-78

CODEN: JIDEAE; ISSN: 0022-202X

DT Journal

LA English

AB Various esters of atropine and scopolamine were most effective in inhibiting sweating after topical application, especially esters of scopolamine-HBr. The effective compds. included esters with straight-chain, branched, cyclic aliphatic, and aromatic groups. Their effectiveness was attributable to their ability to penetrate the skin; as much as 5-10% of the amount applied was absorbed. No systemic after-effects were observed following repeated use of 0.025% solns. over the same area. Benzoylscopolamine did not cause skin irritation or sensitization.

OSC.G 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

L4 ANSWER 12 OF 12 CAPLUS COPYRIGHT 2010 ACS on STN

AN 1963:59688 CAPLUS
 DN 58:59688
 OREF 58:10174d-f
 TI (1-Methyl-2-pyrrolidyl)methyl α -phenyl- α -cyclohexylglycolate
 PA Laboratoires Dausse, S.A.
 SO 14 pp.
 DT Patent
 LA Unavailable

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	FR M1434		19620910	FR	19610719 <--
	FR 1328916			FR	
PRAI	FR		19610719		

GI For diagram(s), see printed CA Issue.

AB HCl gas was passed through 270 g. 1-2-hydroxymethyl-1-methylpyrrolidine in 4.5 l. Et₂O to give the HCl salt (I), SOCl₂ (395 ml.) was added slowly with cooling, the mixture kept 1 hr. at room temperature, and heated 3 hrs. at 100°, 500 ml. C₆H₆ added, the solution evaporated, and the residue washed with C₆H₆ and Et₂O to give 376 g. 1-2-chloromethyl-1-methylpyrrolidine HCl salt, m. 162° (MeCOEt), $[\alpha]_D -5.04 \pm 0.05^\circ$ (50%, H₂O). Alc. KOH (1.66N) (361.3 ml.) was added dropwise to 139 g. dl- α -phenyl- α -cyclohexylglycolic acid in 300 ml. EtOH followed by 102 g. I in 500 ml. EtOH and 361.3 ml. 1.66N alc. KOH, the mixture refluxed 15 min., kept 3 days, filtered, the filtrate evaporated in vacuo, the residue dissolved in 3 l. Et₂O, washed with 10% Na₂CO₃ and H₂O, dried, the solution evaporated, the oil dissolved in 1.5 l. iso-PrOH, HCl gas passed in to pH 3-4, and the solution cooled to give 33% 1-(1-methyl-2-pyrrolidyl)methyl-1- α -phenyl- α -cyclohexylglycolate HCl salt (1-II), m. 220-2°; free base $[\alpha]_D -21 \pm 0.7^\circ$ (4.8, alc.), $-18.4 \pm 0.7^\circ$ (4%, Me₂CO); EtBr salt m. 169-71°, $[\alpha]_D + 10.1 \pm 1^\circ$ (4.3%, alc.). II was similarly prepared in 72% yield from 1- α -phenyl- α -cyclohexylglycolic acid (III). II, $[\alpha]_D 0.9 \pm 0.9^\circ$ (1%, alc.), $-5.2 \pm 1^\circ$ (0.93%, H₂O), was also prepared in 52% yield by refluxing 40 g. III and 23 g. I for 8 hrs. in 250 ml. iso-PrOH and cooling the mixture overnight. II has antispasmodic properties.